

日本における気象衛星利用による文献目録 (1967年1月～1985年4月)

Bibliographical References to Works Dealing with the Use of Satellite Data in Practical Work and Research published in Japan*

- Abe, K., K. Yamamoto, 1979: Determination of sea surface temperature from the Geostationary Meteorological Satellite "Himawari" infrared radiation data, *TENKI*, Vol. 26, No. 9, pp. 493-507 (in Japanese).
- Abe, K., 1980: The accuracy of the precipitable water estimated from the Infrared Radiation data of GMS, *MSC Tech. Note*,** No. 2, pp. 77-93 (in Japanese).
- Abe, K., 1981: Precision of ten-day mean sea surface temperature sensing from the GMS Infrared Radiometer, *MSC Tech. Note*, No. 3, pp. 25-35 (in Japanese).
- Abe, K., 1981: On the improvement of the first guess temperature field for the derivation of ten-day mean sea surface temperatures from GMS infrared images, *MSC Tech. Note*, No. 4, pp. 101-116 (in Japanese).
- Abe, K., R. S. Lourens, 1983: A dust storm over Melbourne, *Weather*, Vol. 38, No. 9, pp. 272-275 (in English).
- Adachi, T. and T. Kasai, 1970: Spectroscopic analysis of photographs taken by NIMBUS II APT system (II) – An improvement in the method of the stereoscopic method, *Journal of the Meteorological Society of Japan*, Vol. 48, No. 3, pp. 234-242 (in English).
- Akaeda, K., T. Takeda, 1984: The utilization of NOAA 7 AVHRR channel 3 data to infer the cloud structure—The behavior of mechanical noise contained in channel 3 data and exclusion of this mechanical noise, *TENKI*, Vol. 31, No. 1, pp. 53-58 (in Japanese).
- Akamatsu, H., 1982: On seiches in Nagasaki-Bay, *Papers in Meteorology and Geophysics*, Vol. 33, No. 2, pp. 95-115 (in Japanese).
- Akiyama, T., 1984: A medium-scale cloud cluster in a Baiu front Part I: Evolution process and fine structure, *Journal of the Meteorological Society of Japan*, Vol. 62, No. 3, pp. 485-504 (in English).
- Akiyama, T., 1984: A medium-scale cloud cluster in a Baiu front Part II: Thermal and kinematic fields and heat budget, *Journal of the Meteorological Society of Japan*, Vol. 62, No. 3, pp. 505-521 (in English).

* 編集者：田中 昭，長谷川秀行，操野年之，前橋紀恵子（気象衛星センター・システム管理課）

** Meteorological Satellite Center Technical Note.

METEOROLOGICAL SATELLITE CENTER TECHNICAL NOTE No. 13. MARCH 1986

- Aoki, T., 1978: On the measured transmission function for the pure rotational band of water vapor, Science Report of Tohoku University, Ser. 5 Geophysics, No. 1, pp. 10-24 (in English).
- Aoki, T., 1979: On the information content of the satellite measured infrared radiation in the atmospheric window region, Journal of the Meteorological Society of Japan, Vol. 57, No. 1, pp. 73-78 (in English).
- Aoki, T., 1980: Statistical determination of clear radiance from cloud-contaminated radiances, Journal of the Meteorological Society of Japan, Vol. 58, No. 6, pp. 528-536 (in English).
- Aoki, T., 1980: A method for matching the HIRS/2 and AVHRR pictures of TIROS-N satellites, MSC Tech. Note, No. 2, pp. 15-26 (in Japanese).
- Aoki, T., 1980: An accurate representation of the transmission functions of the H₂O and CO₂ infrared bands, J. Quant. Spectrosc. Transfer, Vol. 24, pp. 191-202 (in English).
- Aoki, T., S. Nakajima, K. Kato, 1981: Data processing system for TIROS-N satellite series, TENKI, Vol. 28, No. 9, pp. 531-542 (in Japanese).
- Aoki, T., 1982: An improved method to retrieve the clear column radiance from partially cloudy spots of radiometer on board satellite, Journal of the Meteorological Society of Japan, Vol. 60, No. 2, pp. 758-764 (in English).
- Aoki, T., S. Nakajima, K. Kato, 1982: Remote measurements of the sea surface temperature by multichannel observation from orbital satellite, Journal of the Meteorological Society of Japan, Vol. 60, No. 6, pp. 1238-1248 (in English).
- Aoki, T., 1982: Theoretical background of the vertical sounding from TIROS-N satellite series, MSC Tech. Note, No. 5, pp. 25-32 (in English).
- Aoki, T., 1982: Microwave remote sensing, Meteorological Research Note, No. 144, pp. 57-71 (in Japanese).
- Aoki, T., T. Inoue, Estimation of the precipitable water from the IR channel of the geostationary satellite, Remote sensing of environment, No. 12, pp. 219-228 (in English).
- Aoki, T., 1983: A generalized representation of objective analysis, Contributions to atmospheric physics, Vol. 56, No. 4, pp. 464-479 (in English).
- Aoki, T., 1983: Clear radiance retrieval of HIRS channels with the use of AVHRR data, The technical proc. of first international TOVS study conference, Igls, Austria, 29 Aug.-2 Sep. 1983 (in English).
- Arai, K., H. Yamashita, 1981: An attempt of monitoring ground surface temperature form GMS, MSC Tech. Note, No. 3, pp. 1-13 (in Japanese).
- Cadet, D.L. and S.H. Houston, 1984: Precipitable water over Africa and the eastern/central atlantic ocean during

the 1979 summer, Journal of the Meteorological Society of Japan, Vol. 62, No. 5, pp. 761-774 (in English).

Forecast Division, Forecast Dept., JMA, 1982: Verification of very short range forecast technique for precipitation using weather radar, AMeDAS and GMS—A summary of national symposium on forecasting technique held in fiscal 1981, Journal of Meteorological Research, Vol. 34, No. 5, pp. 211-238 (in Japanese).

Fujita, T.T., 1982: Principle of stereoscopic height computation and their applications to stratospheric cirrus over severe thunderstorms, Journal of the Meteorological Society of Japan, Vol. 60, No. 1, pp. 355-368 (in English).

Fukui, T., M. Maeda, 1980: Solar noise interference with GMS down link, MSC Tech. Note, No. 2, pp. 3-14, (in Japanese).

Fukui, T., M. Matsuo, 1981: The influence of the ionospheric scintillation on the GMS communication link, MSC Tech. Note, No. 4, pp. 43-61 (in English).

Fukui, T., K. Tachibana, 1984: Extraordinary phenomenon with communication link, Weather Service Bulletin, Vol. 51, No. 2, pp. 69-86 (in Japanese).

Fuchita, N., 1984: On the relative motion of typhoons—In the case of Typhoon 8305 and Typhoons 8306, 8307, MSC Tech. Note, No. 10, pp. 1-8 (in Japanese).

Funada, H., K. Arai, 1983: GMS-2 observation of volcanic ashes from Mexican Volcano EL Chichon, MSC Tech. Note, No. 7, pp. 13-27 (in Japanese).

Godbole, R.V. and V.R. Murty, 1970: The Indian summer monsoon as seen by weather satellite, Journal of the Meteorological Society of Japan, Vol. 48, No. 4, pp. 360-368 (in English).

Hagiwara, T., 1984: Accuracy of typhoon center determined by GMS imagery, MSC Tech. Note, No. 10, pp. 15-18 (in Japanese).

Hamada, T., 1979: On precision of winds extracted from earth disk images taken from geostationary satellite, MSC Tech. Note, No. 1, pp. 33-52 (in Japanese).

Hamada, T., 1979: Brief description for error estimates of satellite winds derived by CWES system, MSC Tech. Note, No. 1, pp. 79-83 (in English).

Hamada, T., 1980: Wind estimation from images ingested from geostationary meteorological satellite “HIMAWARI”, TENKI, Vol. 27, No. 3, pp. 139-158 (in Japanese).

Hamada, T., 1981: Summary of the international comparison of satellite winds, MSC Tech. Note, No. 4, pp. 117-132 (in Japanese).

METEOROLOGICAL SATELLITE CENTER TECHNICAL NOTE No. 13. MARCH 1986

Hamada, T., 1982: New procedure of height assignment to GMS satellite winds, MSC Tech. Note, No. 5, pp. 91-95 (in English).

Hamada, T., 1982: Representative heights of GMS satellite winds, MSC Tech. Note, No. 6, pp. 35-47 (in English).

Hamada, T., K. Kato, 1984: A new procedure of objective target cloud selection for the GMS cloud wind estimation system and the accuracy of the resultant wind vectors, MSC Tech. Note, No. 9, pp. 27-38 (in Japanese).

Harada, T., 1979: Calculation of sub-cloud point, MSC Tech. Note, No. 1, pp. 53-57 (in Japanese).

Harimaya, T., C. Magono, S. Adachi, K. Hozumi, 1978: An estimation of cloud mass flux based on cloud pictures, Journal of the Meteorological Society of Japan, vol. 56, No. 5, pp. 481-488 (in English).

Hasebe, F., 1982: Observation of stratospheric ozone by the meteorological satellite-Nimbus 4 BUV experiment, TENKI, Vol. 29, No. 9, pp. 905-933 (in Japanese).

Hasegawa, R., K. Ninomiya, 1984: Features of Nagasaki heavy rainstorm of 23 July 1982 as revealed by GMS observation data, TENKI, Vol. 31, No. 9, pp. 565-572 (in Japanese).

Hashimoto, S., 1982: Determination of typhoon center using equivalent black body temperature (T) Distribution Pattern, MSC Tech. Note, No. 6, pp. 11-21 (in Japanese).

Hauze, R.A., 1982: Cloud clusters and large-scale vertical motions in the tropics, Journal of the Meteorological Society of Japan, Vol. 60, No. 1, pp. 396-410 (in English).

Hayashi, R., 1983: Modified methods of estimating space-time spectra from polar-orbiting satellite data, Journal of the Meteorological Society of Japan, Vol. 61, No. 2, pp. 254-262 (in English).

Hayashi, R., 1983: Modified methods of estimating space-time spectra from polar-orbiting satellite data. Part II: The wavenumber transform method, Journal of the Meteorological Society of Japan, Vol. 61, No. 2, pp. 263-268 (in English).

Hayashi, S., A. Watanabe, 1982: Short-term variation of surface oceanographic conditions east off Sanriku region obtained from analysis of NOAA-5, VHRR-IR images, Journal of Meteorological Research, Vol. 34, No. 4, pp. 167-174 (in Japanese).

Hiraki, T., 1980: On the utilization of satellite data for objective analysis and numerical prediction, Technical Note in E.C.C. (J.M.A.), No. 33 (in English).

Hirooka, G., 1983: Partial level down in brightness of geostationary meteorological satellite (GMS) infrared (IR) images caused by the moon, MSC Tech. Note, No. 8, pp. 61-66 (in English).

- Hirota, I., K. Saotome, T. Suzuki and S. Ikeda, 1973: Structure and behavior of the Aleutian anticyclone as revealed by meteorological rocket and satellite observations, Journal of the Meteorological Society of Japan, Vol. 51, No. 5, pp. 353-363 (in English).
- Hozumi, K., C. Magono, 1984: The cloud structure of convergent cloud bands over the Japan sea in winter monsoon period, Journal of the Meteorological Society of Japan, Vol. 62, No. 3, pp. 522-533 (in English).
- Ichiki, A., M. Togashi, A. Uchiyama, 1984: An analysis of the data obtained by simultaneous observation of GMS and GMS-2 - Study of the difference between the brightness temperatures by GMS and those by GMS-2, MSC Tech. Note, No. 10, pp. 19-27 (in Japanese).
- Ichizawa, J., 1983: A history of the satellite wind estimation at MSC and the present accuracy of satellite winds, MSC Tech. Note, No. 8, pp. 85-100 (in Japanese).
- Inoue, T., 1985: On the temperature and effective emissivity determination of semi-transparent cirrus clouds by bi-spectral measurements in the 10 μm window region, Journal of the Meteorological Society of Japan, Vol. 63, No. 1, pp. 88-99 (in English).
- Ishihara, M. and Z. Yanagisawa, 1982: Structure of a tropical squall line observed in the western tropical Pacific during MONEX, Papers in Meteorology and Geophysics, Vol. 33, No. 3, pp. 117-135 (in English).
- Ishizaki, H., M. Endoh and T. Tada, 1983: A trial mapping of sea surface height distribution over the Kuroshio extension region by use of SEASAT-1 altimeter data, Papers in Meteorology and Geophysics, Vol. 33, No. 4, pp. 307-314 (in English).
- Itoh, H., T. Tsuchiya, 1982: Some characteristic features of cloud lines seen in GMS imageries, MSC Tech. Note, No. 5, pp. 33-45 (in English).
- Iwabuchi, T., T. Hoshino, S. Katsuki, H. Fujimura, 1983: VISSR observation of the solar eclipse in rapid-interval limited scan mode, MSC Tech. Note, No. 8, pp. 79-84 (in Japanese).
- Izawa, T. and T. Inoue, 1978: On the NOAA VTPR infrared sensing of the atmosphere and ocean surface during the AMTEX periods, Papers in Meteorology and Geophysics, Vol. 29 No. 2, pp. 35-63 (in English).
- Kamiko, T., 1977: A computer technique for the estimation of cloud height from satellite-viewed shadow, The Geophysical Magazine, Vol. 38, No. 1, pp. 71-84 (in English).
- Kato, K., S. Nakajima, M. Ikawa, 1980: An intercomparison of GMS image data and AMeDAS (Automated Meteorological Data Acquisition System) Rainfall data, MSC Tech. Note, No. 2, pp. 43-59 (in Japanese).
- Kato, K., 1981: Reviews of objective cloud type classification using satellite image data, MSC Tech. Note, No. 3, pp. 37-54 (in Japanese).
- Kato, K., M. Ishikawa, 1982: Objective cloud top-height estimation system for GMS imageries, MSC Tech. Note,

No. 5, pp. 1-24 (in Japanese).

Kato K., 1985: On the abrupt change in the structure of the Baiu front over the China continent in late May of 1979, Journal of the Meteorological Society of Japan, Vol. 63, No. 1, pp. 20-36 (in English).

Kato, M., 1980: Comparison of satellite derived cloud wind and rawin-sonde data, MSC Tech. Note, No. 2, pp. 61-75 (in Japanese).

Kato, M., 1983: A comparison of moisture and temperature fields derived from meteorological satellite vertical sounding (NOAA-TOVS) data with rawinsonde data (Part II), MSC Tech. Note, No. 8, pp. 11-40 (in Japanese).

Kato, M., 1982: Moisture field analysis derived from meteorological satellite vertical sounding (NOAA-TOVS) –Case study–, MSC Tech. Note, No. 6, pp. 23-33 (in Japanese).

Katsuki, S., 1983: On the infrared brightness level increase caused by the sun light during the eclipse period, MSC Tech. Note, No. 8, pp. 101-107 (in Japanese).

Kawamura, R., 1984: Relation between atmospheric circulation and dominant sea surface temperature anomaly patterns in the north pacific during the northern winter, Journal of the Meteorological Society of Japan, Vol. 62, No. 6, pp. 910-916 (in English).

Kikuchi, K. and T. Kasai, 1968: Stereoscopic analysis of photographs taken by Nimbus II APT system, Journal of the Meteorological Society of Japan, Vol. 46, No. 1, pp. 60-67 (in English).

Kimura, T., 1983: Analysis of sea ice in the sea of Okhotsk using LANDSAT-MSS data – Comparison with GMS data –, MSC Tech. Note, No. 8, pp. 41-59 (in Japanese).

Kimura, T., 1984: Clouds with appearance like Cb-cloud on satellite imagery, MSC Tech. Note, No. 9, pp. 11-26 (in Japanese).

Koba, H., 1983: Features of satellite observed cloud-top temperature contour pattern of convective cloud clusters with heavy rainfall in Baiu-season – Case study –, MSC Tech. Note, No. 7, pp. 1-12 (in Japanese).

Koba H., 1984: An explanation of the revised DVORAK technique for estimating tropical cyclone intensity using satellite cloud imagry and some results of its application to two 1982 storms, MSC Tech. Note, No. 9, pp. 39-56 (in Japanese).

Kodaira, N., N. Murayama, Y. Takayama and E. Kamishiro, 1976: The data collection platform of the GMS system, Papers in Meteorology and Geophysics, Vol. 27, No. 3, pp. 63-73 (in English).

Kodaira, N., K. Kato, T. Hamada, 1979: Man-machine interactive processing of extracting cloud top height and cloud wind data from the GMS images, MSC Tech. Note, No. 1, pp. 59-78 (in English).

- Kodaira, N., N. Murayama, 1982: Meteorological observation using remote sensing technique, TENKI, Vol. 29, No. 2, pp. 113-133 (in Japanese).
- Konage, S. and H. Ishizaki, 1981: Observation of short-term sea-level fluctuation around the Kuroshio south of Honshu, Japan by the Seasat-a altimetry, Papers in Meteorology and Geophysics, Vol. 32, No. 3, pp. 183-192 (in English).
- Kodaira, N., N. Murayama, 1982: Meteorological observation using remote sensing technique, TENKI, Vol. 29, No. 2, pp. 113-133 (in Japanese).
- Konaga, S. and H. Ishizaki, 1981: Observation of short-term sea-level fluctuation around the Kuroshio south of Honshu, Japan by the Seasat-a altimetry, Papers in Meteorology and Geophysics, Vol. 32, No. 3, pp. 183-192 (in English). Observation of short-term sea-level fluctuation around the Kuroshio south of Honshu, Japan by the Seasat-a altimetry, Papers in Meteorology and Geophysics, Vol. 32, No. 3, pp. 183-192 (in English). Observation of short-term sea-level fluctuation around the Kuroshio south of Honshu, Japan by the Seasat-a altimetry, Papers in Meteorology and Geophysics, Vol. 32, No. 3, pp. 183-192 (in English).
- Kuroda, Y., 1981: Convective cloud band associated with easterly wave, MSC Tech. Note, No. 4, pp. 21-42 (in Japanese).
- Kuroda, Y., 1982: Characteristic cloud patterns associated with the heavy rainfall occurred over the Hokkaido district 3 to 6 August 1981, MSC Tech. Note, No. 6, pp. 73-90 (in Japanese).
- Love, G., 1985: A study of the linear relationships between the monthly mean fields of sea surface temperature, mean sea level pressure and cloudiness over the northwest pacific, Journal of the Meteorological Society of Japan, Vol. 63, No. 2, pp. 201-209 (in English).
- Maeda, N., T. Takahashi, 1984: VISSR IR grid data processing, MSC Tech. Note, No. 9, pp. 57-60 (in Japanese).
- Murayama, T., 1982: Upper tropospheric zonal wind asciliation with a 30-50 day period over the equatorial western Pacific observed in cloud movement vectors, Journal of the Meteorological Society of Japan, Vol. 60, No. 1, pp. 172-182 (in English).
- Matsumoto, S., K. Ninomiya, K. Hasegawa and Y. Miki, 1982: The structure and the role of a subsynoptic-scale cold vortex on the heavy precipitation, Journal of the Meteorological Society of Japan, Vol. 60, No. 1, pp. 339-354 (in English).
- Matsumoto, S., H. Okamura, 1985: The interval gravity wave observed in the typhoon T8124 (Gay), Journal of the Meteorological Society of Japan, Vol. 63, No. 1, pp. 37-51 (in English).
- Matsunaga, T., 1983: The isothermal pattern mapping of the GMS VISSR infrared imagery and its application to a heavy rainstorm, MSC Tech. Note, No. 7, pp. 39-46 (in English).
- Matsuzaki, A., T. Itoh and Y. Nakamura, 1984: On observation of middle atmosphere with LAS (Limb-atmospheric Infrared Spectrometer) on board of satellite "Ohzora" (EXOS-C), Journal of the Meteorological Society of Japan, Vol. 62, No. 2, pp. 328-333 (in English).

METEOROLOGICAL SATELLITE CENTER TECHNICAL NOTE No. 13. MARCH 1986

- McFarland, M.J., Y.K. Sasaki, 1977: Variational analysis of temperature and moisture advection in a severe storm environment, Journal of the Meteorological Society of Japan, Vol. 55, No. 4, pp. 421-430 (in English).
- Mitate, Y., M. Hitsuma, 1984: A supplement study on the subsynoptic cloud system off the South Coast of Japan, MSC Tech. Note, No. 10, pp. 57-60 (in Japanese).
- Miura, N., H. Ohno, 1983: A new display technique (satellite imagery oriented) for the chart of predicted overcast area, TENKI, Vol. 30, No. 6, pp. 303-309 (in Japanese).
- Momose, H., 1975: Analysis of cloud amount over the China continent and the East China Sea, Journal of the Meteorological Society of Japan, Vol. 53, No. 1, pp. 87-91 (in English).
- Morikawa, M., 1982: Extraction of VISSR image data with a polygonal shape, MSC Tech. Note, No. 6, pp. 49-58 (in Japanese).
- Moriwaki, A., 1984: The Karman vortex street detected by the Geostationary meteorological satellite "Himawari", MSC Tech. Note, No. 9, pp. 1-9 (in Japanese).
- Motoki, T., M. Satomi, 1981: Short-interval observation of winter monsoon cloud pattern by GMS-1 (part-1), MSC Tech. Note, No. 3, pp. 15-23 (in Japanese).
- Motoki, T., A. Kamoshida, Y. Shimizu, T. Tokita, 1983: Outline of operational production of T_{BB} contour map and characteristics of the computational technique, Weather Service Bulletin, Vol. 50, No. 6, pp. 461-469 (in Japanese).
- Motoki, T., A. Kamoshida, 1984: Analysis of T_{BB} contour map, Weather Service Bulletin, Vol. 51, No. 2, pp. 61-67 (in Japanese).
- Motoki, T., H. Tani, I. Horiguchi, 1984: Application of satellite data to the studies of agricultural meteorology – Relationship between ground temperature from GMS IR data and AMeDAS air temperature –, Journal of Agricultural Meteorology, Vol. 40, No. 2, pp. 111-117 (in Japanese).
- Murakami, T., F.P. Ho, 1972: Spectrum analysis of cloudiness over the Northern Pacific, Journal of the Meteorological Society of Japan, Vol. 50, No. 4, pp. 285-311 (in English).
- Murakami, T., 1976: Cloudiness fluctuations during the summer monsoon, Journal of the Meteorological Society of Japan, Vol. 54, No. 3, pp. 175-181 (in English).
- Muramatsu, T., 1980: The cloud line enhanced by upwind orographic features winter monsoon situation, Geographical Magazine, Vol. 38, No. 4, pp. 15 (in English).
- Muramatsu, T., 1983: Diurnal variations of satellite-measured T_{BB} areal distribution and eye diameter of mature typhoons, Journal of the Meteorological Society of Japan, Vol. 61, No. 1, pp. 77-90 (in English).

- Nagai, T., 1984: On the P1 data from GMS/SEM, MSC Tech. Note, No. 10, pp. 37-46 (in Japanese).
- Nagasaki, K., 1982: Preliminary evaluation of sea surface temperature derived from IR observations of the GMS, The Oceanographical Magazine, Vol. 32, pp. 19-24 (in English).
- Nakamura, K., K. Sugimoto, 1980: On the program to print out the distribution of the brightness temperature and albedo value of the VISSR data and its application, Weather Service Bulletin, Vol. 47, No. 1-2, pp. 7-17 (in Japanese).
- Nakamura, K., T. Tsuchiya, 1980: An examination of the reliability of densitometric analysis, MSC Tech. Note, No. 2, pp. 95-103 (in English).
- Nakamura, K., K. Sugimoto, 1981: On the development of image data edition and display system – For the determination of the cloud system center of a typhoon and the analysis of a meso-scale phenomenon –, MSC Tech. Note, No. 4, pp. 141-154 (in Japanese).
- Nasuda, H., 1985: Infrared greybody emissivity and visible albedo of high altitude semitransparent cloud, Journal of the Meteorological Society of Japan, Vol. 63, No. 1, pp. 75-87 (in English).
- Ninomiya, K., T. Akiyama, 1973: Satellite and radar observational study of the wave disturbances over the east coast of the Asian continent in February 1968, Journal of the Meteorological Society of Japan, Vol. 51, No. 6, pp. 497-500 (in English).
- Ninomiya, K., T. Akiyama, 1981: Long-lived medium-scale cumulonimbus cluster in Asian subtropic humid region, Journal of the Meteorological Society of Japan, Vol. 59, No. 4, pp. 564-577 (in English).
- Ninomiya, K., R. Hasegawa, Y. Tatsumi, 1983: Subtropical rainband along the western side of a slow-moving ridge – Observation and real data forecast experiment, Journal of the Meteorological Society of Japan, Vol. 61, No. 4, pp. 606-618 (in English).
- Nirei, S., 1981: Comparison of the cloud top heights evaluated from the GMS image with the heights reported by pilots, MSC Tech. Note, No. 4, pp. 69-100 (in Japanese).
- Nirei, S., 1983: Comparison of the cloud top heights evaluated from the GMS infrared image with the heights reported by air-line pilots. (part-2), MSC Tech. Note, No. 8, pp. 1-9 (in English).
- Nitta, T. and K. Masuda, 1981: Observational study of a monsoon depression developed over the bay of Bengal during summer MONEX, Journal of the Meteorological Society of Japan, Vol. 59, No. 5, pp. 672-682 (in English).
- Nitta, T., Y. Nakagomi, Y. Suzuki, N. Hasegawa and A. Kadokura, 1985: Global analysis of lower tropospheric disturbances in the tropics during the northern summer of the FGGE year. Part I: Global features of the disturbances, Journal of the Meteorological Society of Japan, Vol. 63, No. 1, pp. 1-99 (in English).

METEOROLOGICAL SATELLITE CENTER TECHNICAL NOTE No. 13. MARCH 1986

- Ohta, H., 1983: Rapid fading of the signal level on 2 GHz microwave communication link between CDAS and MSC, MSC Tech. Note, No. 8, pp. 67-78 (in Japanese).
- Okabayashi, T., 1982: The use of meteorological satellite data (I), Weather Service Bulletin, Vol. 49, No. 1, pp. 1-50 (in Japanese).
- Okabayashi, T., 1982: The use of meteorological satellite data (II), Weather Service Bulletin, Vol. 49, No. 4, pp. 185-250 (in Japanese).
- Onoda, H., 1980: Cloud distribution obtained by GMS data, TENKI, Vol. 27, No. 6, pp. 395-404 (in Japanese).
- Okamura, H. and I. Isozaki, 1983: Observation of a surface wind around a typhoon by Seasat-1 scatterometer, Papers in Meteorology and Geophysics, Vol. 34, No. 2, pp. 83-94 (in Japanese).
- Orlanski, I. and L.J. Polinsky, 1977: Spectral distribution of cloud cover over Africa, Journal of the Meteorological Society of Japan, Vol. 55, No. 5, pp. 483-493 (in English).
- Ozawa, T., 1977: A case study of the medium scale disturbance: Tosaoki Low, Papers in Meteorology and Geophysics, Vol. 28, No. 2, pp. 63-81 (in Japanese).
- Sakakibara, H., 1981: Heavy rainfall from very shallow convective clouds, Journal of the Meteorological Society of Japan, Vol. 59, No. 3, pp. 387-394 (in English).
- Sakakibara, H., 1983: Meso-and small-scale structures of a cold-frontal precipitation system affected by a small low and mesoscale orography, Papers in Meteorology and Geophysics, Vol. 34, No. 2, pp. 61-73 (in English).
- Sasaki, H., 1985: Clear sky brightness temperature over the Western North Pacific Ocean from GMS Spacecraft, MSC Tech. Note. No. 11, pp. 11-22 (in Japanese).
- Sawada, Y., 1983: Analysis of eruption clouds by the 1981 eruptions of Alaid and Pagan volcanoes with GMS images, Papers in Meteorology and Geophysics, Vol. 34, No. 4, pp. 307-324 (in Japanese).
- Sekihsara, K., Y. Suzuki, and K. Murai, 1978: On the formation of convective cloud cells during cold air outbreaks in AMTEX '75, Papers in Meteorology and Geophysics, Vol. 29, No. 2, pp. 65-73 (in English).
- Sekioka, M., 1970: On the behavior of cloud patterns as seen on satellite photographs in the transformation of a typhoon into an extratropical cyclone, Journal of the Meteorological Society of Japan, Vol. 48, No. 3, pp. 224-233 (in English).
- Serebreny, S.M. and Jr. R.H. Blackmer, 1968: Relationships between radon-detected maritime precipitation and satellite, viewed cloud cover, Journal of the Meteorological Society of Japan, Vol. 46, No. 2, pp. 133-149 (in English).

- Shimada, K., 1982: Prediction of tropical cyclone intensity using equivalent black body temperatures (T_{BB}) measured by GMS, MSC Tech. Note, No. 6, pp. 1-10 (in Japanese).
- Shimada, K., 1984: On the error of typhoon intensity estimation by means of IR dvorak technique occurring when the eye is very small, MSC Tech. Note, No. 10, pp. 9-13 (in Japanese).
- Shimamura, M., 1980: The application of meteorological satellite data in analysis of convective clouds and disturbances in tropics, TENKI, Vol. 27, No. 4, pp. 245-259 (in Japanese).
- Shimamura, M., 1982: An application of GMS satellite data in the analysis of the Upper Cold Low in the Western North Pacific, The Geophysical Magazine, Vol. 40, No. 2, pp. 113-152 (in English).
- Shimamura, M. and H. Yamashita, 1983: An application of satellite data in the typhoon forecast in middle latitudes, The Geophysical Magazine, Vol. 40, No. 4, pp. 313-336 (in English).
- Shimizu, N., 1981: Cloud activity and center pressure of a typhoon, Geophysical Magazine, Vol. 39, No. 4, pp. 267-276 (in English).
- Sikdar, D.N. and Cram, R.S., 1974: Upper tropospheric isobaric temperature and height fields over the tropical atlantic from satellite infrared spectrometer soundings, Journal of the Meteorological Society of Japan, Vol. 52, No. 2, pp. 175-187 (in English).
- Sugi, M. and M. Kanamitsu, 1982: A study of a subtropical upper level cyclone using JMA operational forecast model, Journal of the Meteorological Society of Japan, Vol. 60, No. 4, pp. 932-946 (in English).
- Sugiura, S., 1983: A study of typhoons using SASS wind data observed by Seasat-1, Papers in Meteorology and Geophysics, Vol. 34, No. 1, pp. 35-52 (in Japanese).
- Sumi, A. and T. Murakami, 1981: Large-scale aspects of the 1978-79 winter circulation over the greater WMONEX region, Part I: Monthly and season mean fields, Journal of the Meteorological Society of Japan, Vol. 59, No. 5, pp. 625-645 (in English).
- Sumi, A., 1982: On an evaluation of data impact with respect to Rossby mode in the data assimilation cycle, Journal of the Meteorological Society of Japan, Vol. 60, No. 4, pp. 917-931 (in English).
- Suzuki, K., 1980: Evaluation of cloud top height from infrared radiation data of the Geostationary Meteorological Satellite "HIMAWARI", TENKI, Vol. 27, No. 5, pp. 307-320 (in Japanese).
- Suzuki, K., 1985: Occurrence of linearly ranged Cb-clouds associated with migrating band-shaped cirriform-clouds - A case study -, MSC Tech. Note, No. 11, pp. 1-9 (in Japanese).
- Tada, K., 1981: An atmospheric moisture retrieval technique with radiance data from the satellite, The Geophysical Magazine, Vol. 39, pp. 249-265 (in English).

METEOROLOGICAL SATELLITE CENTER TECHNICAL NOTE No. 13. MARCH 1986

- Tada K., 1981: Use of satellite moisture information from TIROS-N in numerical weather prediction. The Geophysical Magazine, Vol. 40, No. 1, pp. 1-29 (in English).
- Takahashi, T., 1979: The simplified mapping equation of VISSR image data from the Geostationary Meteorological Satellite (GMS), MSC Tech. Note, No. 1, pp. 13-19 (in English).
- Takashima, T. and Y. Takayama, 1981: Estimation of sea surface temperature from remote sensing in the 3.7 μ m window region, Journal of the Meteorological Society of Japan, Vol. 59, No. 6, pp. 876-891 (in English).
- Takashima, T. and Y. Takayama, 1980: A feasibility study to derive the sea surface wind by the use of the sun glint pattern, Papers in Meteorology and Geophysics, Vol. 31, No. 3/4, pp. 173-176 (in English).
- Takashima, T., Y. Takayama, K. Matsuura and K. Naito, 1982: Cloud height determination by satellite stereoscopy, Papers in Meteorology and Geophysics, Vol. 33, No. 2, pp. 65-78 (in English).
- Takeda, A., 1980: Remote sensing of the winds over the oceans from a satellite-Technique and theory of Seasat-A Scatterometer System, TENKI, Vol. 27, No. 2, pp. 87-102 (in Japanese).
- Takeda, T. and M. Ikeyama, 1984: Time variation of cloud amount with about 30-days period in the western north pacific region, Journal of the Meteorological Society of Japan, Vol. 62, No. 4, pp. 165-171 (in English).
- Takeuchi, M. and T. Uchiyama, 1985: Synoptic analysis of polar lows over the Japan sea area and heavy snowfalls in the coastal plain regions, Papers in Meteorology and Geophysics, Vol. 36, No. 1, pp. 1-21 (in Japanese).
- Tanaka, H. and O. Ryuguji, 1971: Spectrum analysis of tropical cloudiness (I), Journal of the Meteorological Society of Japan, Vol. 49, No. 1, pp. 13-19 (in English).
- Tanaka, H. and O. Ryuguji, 1973: Spectrum analysis of tropical cloudiness (II), Journal of the Meteorological Society of Japan, Vol. 51, No. 2, pp. 93-100 (in English).
- Tanaka, Y., K. Tsuchiya and U. Yamaura, 1981: Detection of volcanic smoke and ash-fall area at volcano Aso, from Landsat MSS data, Papers in Meteorology and Geophysics, Vol. 32, No. 4, pp. 275-290 (in English).
- Tik, Chiyu, 1984: A case study of heavy rain spell on 13th-25th December 1982 over the east coast of Peninsular Malaysia and Singapore, Journal of the Meteorological Society of Japan, Vol. 62, No. 2, pp. 296-307 (in English).
- Togashi, M., A. Ichiki, T. Takahasi and S. Katsuki, 1985: Simultaneous observation of GMS and GMS-2 -A study of the difference between the brightness temperatures by GMS and those by GMS-2-, MSC Tech. Note, No. 11, pp. 23-32 (in Japanese).
- Tokuno, M., 1984: The development of image processing of NOAA AVHRR data and its application to sea

surface temperature, MSC Tech. Note, No. 10, pp. 47-55 (in Japanese).

Tsuchiya, K. and T. Fujita, 1967: A satellite meteorological study of evaporation and cloud formation over the western pacific under the influence of the winter monsoon, Journal of the Meteorological society of Japan, Vol. 45, No. 3, pp. 232-250 (in English).

Tsuchiya, T., 1981: Detection of high probable areas of turbulence on satellite imageries; MSC Tech. Note, No. 4, pp. 1-20 (in English).

Tsuchiya, T., R. Ito and C. Ishida, Characteristics of the detectors of multi spectral scanner (MSS) of Landsat in space environment, Journal of the Meteorological Society of Japan, Vol. 60, No. 5, pp. 1165-1174 (in English).

Tsuchiya, T., 1984: Rainfall estimation from meteorological satellite data, TENKI, Vol. 31, No. 10, pp. 607-623 (in Japanese)

Typhoon research division, MRI, 1985: A study on the changes of the three-dimensional structure and the movement speed of the typhoon through its life time, Technical reports of the Meteorological Research Institute, No. 14 (in Japanese).

Uchida, M., 1979: Correspondence of Satellite Cloud Pictures to Charts of Vorticity, Vertical Velocity and Moisture, Journal of Meteorological Research, Vol. 31, No. 3-4, pp. 73-77 (in Japanese).

Uchiyama, A., 1984: VISSR infrared calibration by means of calculated radiance, MSC Tech. Note, No. 10, pp. 29-36 (in Japanese).

Ueda, S., 1983: Ionospheric scintillation which disturbed the GMS-2 communication link, MSC Tech. Note, No. 7, pp. 29-38 (in Japanese).

Watanabe, Y., 1979: V. Dvorak and S. Wright; Tropical cyclone intensity analysis using enhanced IR imagery, Weather Service Bulletin, Vol. 46, No. 9-10, pp. 279-288 (in Japanese).

Yagi, S., Y. Uchiyama, 1983: Cloud movement and confluence around Noto Peninsula and Sado Island in relation to heavy snowfall over Johetsu area, TENKI, Vol. 30, No. 6, pp. 291-294 (in Japanese).

Yamashita, H., 1979: The effects of different time intervals on the wind estimation, MSC Tech. Note, No. 1, pp. 1-11 (in English).

Yasoshina, H., 1982: The GMS Satellite Observation of Volcanic Eruptions in 1981 in the Western Pacific, MSC Tech. Note, No. 5, pp. 47-64 (in Japanese).

Yasunari, T., 1977: Stationary waves in the southern hemisphere mid-latitude zone revealed from average brightness charts, Journal of the Meteorological Society of Japan, Vol. 55, No. 3, pp. 274-285 (in English).

METEOROLOGICAL SATELLITE CENTER TECHNICAL NOTE No. 13. MARCH 1986

Yasunari, T., 1979: Cloudiness fluctuations associated with the northern hemisphere summer monsoon, Journal of the Meteorological Society of Japan, Vol. 57, No. 3, pp. 227-242 (in English).